Brain Research, 417 (1987) 409–411
Elsevier

Author Index

Abood, M.E., Lee, N.M. and Loh, H.H., Modification of opioid agonist binding by pertussis toxin, 70
Aguila, M.C., Khorram, O. and McCann, S.M., α-Melanocyte stimulating hormone discloses a stimulatory effect of β-endorphin on somatostatin release, 127
Akaike, N., see Oyama, Y., 143
Akaike, T., Electrophysiological analysis of the tecto-olivo-cerebellar (lobus simplex) projecion in the rat, 371
Avoli, M., Louvel, J., Pumain, R. and Olivier, A., Seizure-like discharges induced by lowering $[Mg^{2+}]_o$ in the human epileptogenic neocortex maintained in vitro, 199

Baba, S., see Inui, A., 355
Beal, M.F., see Ellison, D.W., 389
Bean, A.J., see Deutch, A.Y., 350
Bell Jr., C.E., see Frikke, M.J., 283
Bell, G.A., see Slotnick, B.M., 343
Benowitz, L.I. and Schmidt, J.T., Activity-dependent sharpening of the regenerating retinotectal projection in goldfish: relationship to the expression of growth-associated proteins, 118
Berdan, R.C. and Bulloch, A.G.M., Tannic acid enhances staining of microtubule associated proteins, but impairs neuronal physiology, 153
Berger, A.J., see Holtman Jr., J.R., 12
Bissette, G., see Deutch, A.Y., 350
Bland, B.H., see Konopacki, J., 399
Boegman, R.J., Metcalf, R., Riopelle, R.J. and Ludwin, S.K., Neurotoxicity of quinolinate in the rat nucleus basalis magnocellularis, 315
Buck, S.H., see Swenberg, M.-L., 131
Bulloch, A.G.M., see Berdan, R.C., 153

Cadete-Leite, A., see Paula-Barbosa, M., 139
Caspary, D.M., Pazara, K.E., Kössl, M. and Faingold, C.L., Strychnine alters the fusiform cell ouput from the dorsal cochlear nucleus, 273
Cervera, P., see Mourre, C., 21
Chai, C.Y., see Kuo, J.S., 181
Chambers, J.P., see Vaughan, M.K., 321
Chang, Y.T., Lin, J.W. and Faber, D.S., Spinal inputs to the ventral dendrite of the teleost Mauthner cell, 205
Chiang, B.N., see Kuo, J.S., 181
Christakos, C.N., see Cohen, M.I., 148
Chung, J.M., see Kim, J., 304
Clarke, D.W., see Masters, B.A., 247
Cohen, M.I., See, W.R., Christakos, C.N. and Sica, A.L., High-frequency and medium-frequency components of different inspiratory nerve discharges and their modification by various inputs, 148

Decker, M.W. and Gallagher, M., Scopolamine-disruption of radial arm maze performance: modification by noradrenergic depletion, 59
Deutch, A.Y., Bean, A.J., Bissette, G., Nemeroff, C.B., Robbins, R.J. and Roth, R.H., Stress-induced alterations in neurotensin, somatostatin and corticotropin-releasing factor in mesotelencephalic dopamine system regions, 350
Dick, T.E., see Holtman Jr., J.R., 12
Dragunow, M. and Robertson, H.A., 8-Cyclopentyl 1,3-dimethylxanthine prolongs epileptic seizures in rats, 377
Drescher, D.G., see Drescher, M.J., 39
Drescher, M.J., Drescher, D.G. and Hatfield, J.S., Potassium-evoked release of endogenous primary amine-containing compounds from the trout saccular macula and saccular nerve in vitro, 39

Ekman, R., see Wahlestedt, C., 33
Ellison, D.W., Beal, M.F. and Martin, J.B., Phosphoethanolamine and ethanolamine are decreased in Alzheimer's disease and Huntington's disease, 389
Espinoza, S.G., see Thomas, H.C., 214

Faber, D.S., see Chang, Y.T., 205
Faingold, C.L., see Caspary, D.M., 273
Ferino, F., Thierry, A.M., Saffroy, M. and Glowinski, J., Interhemispheric and subcortical collaterals of medial prefrontal cortical neurons in the cat, 257
Fosset, M., see Lombet, A., 327
Frikke, M.J., Seshi, B. and Bell Jr., C.E., Monoclonal antibodies to human neuron-specific enolase reveal heterogeneity of the enzyme in neurons of the central nervous system, 283
Fukuda, H., see Kaneko, T., 403

Gallagher, J.P., see Joels, M., 99
Gallagher, M., see Decker, M.W., 59
Gash, D.M., see Kordower, J.H., 85
Glass, E.J., see Hsu, L.L., 232
Glowinski, J., see Ferino, F., 257
Graham, S., see Slotnick, B.M., 343

Hahn, M.E., see Hewitt, J.K., 225
Håkanson, R., see Wahlestedt, C., 33
Hartung, H.-P. and Toyka, K.V., Phorbol diester TPA elicits prostaglandin E release from cultured rat astrocytes, 347
Hatfield, J.S., see Drescher, M.J., 39
Heilig, M., see Wahlestedt, C., 33
Hewitt, J.K., Hahn, M.E. and Karkowski, L.M., Genetic selection disrupts stability of mouse brain weight development, 225
Hodin, R.A., see Kreisman, N.R., 335
Hoffmann, K., see Illnerová, H., 367
Holdefer, R.N. and Jensen, R.A., The effects of peripheral D-amphetamine, 4-OH amphetamine, and epinephrine on maintained discharges in the locus coeruleus with reference to the modulation of learning and memory by these substances, 108
Holtman Jr., J.R., Dick, T.E. and Berger, A.J., Serotonin-mediated excitation of recurrent laryngeal and phrenic motoneurons evoked by stimulation of the raphe obscurus, 12
Hori, N., see Oyama, Y., 143
Hsu, L.L., Yu, J.-R., Upp Jr., J.R., Glass, E.J. and Townsend, C.M., Chronic bombesin treatment increased the [^{3}H]-

spiperone binding, glutamate decarboxylase and choline acetyltransferase activity in the rat brain, 232
Huston, J.P., see Schwarting, R., 75

Igarashi, M., see Usami, S., 367
Illnerová, H., Vanecek, J. and Hoffmann, K., Adjustment of the rat pineal N-acetyltransferase rhythm to eight-hour shifts of the light-dark cycle: advance of the cycle disturbs the rhythm more than delay, 367
Inoue, T., see Inui, A., 355
Inoue, Y., see Terashima, T., 190
Inui, A., Inoue, T., Sakatani, N., Oya, M., Morioka, H. and Baba, S., Proglumide has access to brain and antagonizes the central satiety effect of cholecystokinin octapeptide in the dog, 355

Jacomet, Y., see Lombet, A., 327
Jensen, R.A., see Holdefer, R.N., 108
Joels, M., Shinnick-Gallagher, P. and Gallagher, J.P., Effect of serotonin and serotonin analogues on passive membrane properties of lateral septal neurons in vitro, 99
Jourdan, F., see Royet, J.P., 1
Judkins, J.H., see Masters, B.A., 247

Kalivas, P.W., see Vezina, P., 51
Kaneko, T., Ono, H. and Fukuda, H., Enhancement of recurrent inhibition of the spinal monosynaptic reflex by preceding stimulation of the medullary raphé in rats, 403
Karkowski, L.M., see Hewitt, J.K., 225
Katada, T., see Terashima, T., 190
Kehr, A.D., see Mufson, E.J., 385
Kennedy, P.R., Parametric relationships of individual digit movements to neuronal discharges in primate magnocellular red nucleus, 185
Khorram, O., see Aguila, M.C., 127
Kim, J., Shin, H.K. and Chung, J.M., Many ventral root afferent fibers in the cat are third branches of dorsal root ganglion cells, 304
Konopacki, J., Bland, B.H. and Roth, S.H., Phase shifting of CA_1 and dentate EEG theta rhythms in hippocampal formation slices, 399
Kordower, J.H., Notter, M.F.D. and Gash, D.M., Neuroblastoma cells in neural transplants: a neuroanatomical and behavioral analysis, 85
Kössl, M., see Caspary, D.M., 273
Kreisman, N.R., Hodin, R.A., Rosenthal, M. and Sick, T.J., Role of pulmonary edema in phasic changes of cerebral oxygenation during serial seizures, 335
Kuo, J.S., Wang, M.R., Liu, R.H., Yu, C.Y., Chiang, B.N. and Chai, C.Y., Reduction of common carotid resistance upon stimulation of an area dorsal to the facial nucleus of cats, 181

Laing, D.G., see Slotnick, B.M., 343
Lazdunski, M., see Lombet, A., 327
Lazdunski, M., see Mourre, C., 21
Lee, N.M., see Abood, M.E., 70
Leon, M., see Wilson, D.A., 175
Leonard, C.M., see Wirsig, C.R., 293
Le Roith, D., see Masters, B.A., 247
Lieberburg, I., Developmental expression and regional distribution of the scrapie-associated protein and mRNA in the rat central nervous system, 363
Lin, J.W., see Chang, Y.T., 205

Liu, R.H., see Kuo, J.S., 181
Loh, H.H., see Abood, M.E., 70
Lombet, A., Fosset, M., Romey, G., Jacomet, Y. and Lazdunski, M., Identification in mammalian brain of an endogenous substance with Na^+ channel blocking activities similar to those of tetrodotoxin, 327
Louvel, J., see Avoli, M., 199
Lovenberg, W., see Swenberg, M.-L., 131
Ludwin, S.K., see Boegman, R.J., 315

Martin, J.B., see Ellison, D.W., 389
Masters, B.A., Shemer, J., Judkins, J.H., Clarke, D.W., Le Roith, D. and Raizada, M.K., Insulin receptors and insulin action in dissociated brain cells, 247
Matsui, H., Effect of subthalamic stimulation on adrenal epinephrine and norepinephrine secretion in the rat, 158
McCann, S.M., see Aguila, M.C., 127
McCaslin, P.P. and Morgan, W.W., Cultured cerebellar cells as an in vitro model of excitatory amino acid receptor function, 380
Mesulam, M.-M., see Mufson, E.J., 385
Metcalf, R., see Boegman, R.J., 315
Mink, J.W. and Thach, W.T., Preferential relation of pallidal neurons to ballistic movements, 393
Morgan, W.W., see McCaslin, P.P., 380
Morioka, H., see Inui, A., 355
Mourre, C., Cervera, P. and Lazdunski, M., Autoradiographic analysis in rat brain of the postnatal ontogeny of voltage-dependent Na^+ channels, Ca^{2+}-dependent K^+ channels and slow Ca^{2+} channels identified as receptors for tetrodotoxin, apamin and (–)-desmethyoxyverapamil, 21
Mufson, E.J., Kehr, A.D., Wainer, B.H. and Mesulam, M.-M., Cortical effects of neurotoxic damage to the nucleus basalis in rats: persistent loss of extrinsic cholinergic input and lack of transsynaptic effect upon the number of somatostatin-containing, cholinesterase-positive, and cholinergic cortical neurons, 385

Nagel, J.A., see Schwarting, R., 75
Nakajima, T., Yashima, Y. and Nakamura, K., Higher density of ^{125}I-neuropeptide Y receptors in the area postrema of SHR, 360
Nakamura, K., see Nakajima, T., 360
Nemeroff, C.B., see Deutch, A.Y., 350
Notter, M.F.D., see Kordower, J.H., 85

Oinuma, M., see Terashima, T., 190
Okada, Y., see Tanimoto, M., 239
Olivier, A., see Avoli, M., 199
Ono, H., see Kaneko, T., 403
Osmanović, S.S. and Shefner, S.A., Anomalous rectification in rat locus coeruleus neurons, 161
Oya, M., see Inui, A., 355
Oyama, Y., Hori, N., Tokutomi, N. and Akaike, N., D-600 blocks open Ca^{2+} channels more profoundly than closed ones, 143

Paula-Barbosa, M., Tavares, M.A. and Cadete-Leite, A., A quantitative study of frontal cortex dendritic microtubules in patients with Alzheimer's disease, 139
Pazara, K.E., see Caspary, D.M., 273
Pumain, R., see Avoli, M., 199

Raizada, M.K., see Masters, B.A., 247

Reiter, R.J., see Vaughan, M.K., 321
Riopelle, R.J., see Boegman, R.J., 315
Robbins, R.J., see Deutch, A.Y., 350
Robertson, H.A., see Dragunow, M., 377
Romey, G., see Lombet, A., 327
Rosenthal, M., see Kreisman, N.T., 335
Roth, R.H., see Deutch, A.Y., 350
Roth, S.H., see Konopacki, J., 399
Royet, J.P., Sicard, G., Souchier, C. and Jourdan, F., Specificity of spatial patterns of glomerular activation in the mouse olfactory bulb: computer-assisted image analysis of 2-deoxyglucose autoradiograms, 1

Saffroy, M., see Ferino, F., 257
Sagar, S.M., Sharp, F.R. and Swanson, R.A., The regional distribution of glycogen in rat brain fixed by microwave irradiation, 172
Saito, Y. and Wright, E.M., Regulation of intracellular chloride in bullfrog choroid plexus, 267
Sakatani, N., see Inui, A., 355
Schmidt, J.T., see Benowitz, L.I., 118
Schwarting, R., Nagel, J.A. and Huston, J.P., Asymmetries of brain dopamine metabolism related to conditioned paw usage in the rat, 75
See, W.R., see Cohen, M.I., 148
Seshi, B., see Frikke, M.J., 283
Sharp, F.R., see Sagar, S.M., 172
Shefner, S.A., see Osmanovic, S.S., 161
Shemer, J., see Masters, B.A., 247
Shin, H.K., see Kim, J., 304
Shinnick-Gallagher, P., see Joels, M., 99
Sica, A.L., see Cohen, M.I., 148
Sicard, G., see Royet, J.P., 1
Sick, T.J., see Kreisman, N.R., 335
Skagerberg, G., see Wahlestedt, C., 33
Slotnick, B.M., Graham, S., Laing, D.G. and Bell, G.A., Detection of propionic acid vapor by rats with lesions of olfactory bulb areas associated with high 2-DG uptake, 343
Souchier, C., see Royet, J.P., 1
Stewart, J., see Vezina, P., 51
Sundler, F., see Wahlestedt, C., 33
Swanson, R.A., see Sagar, S.M., 172
Swenberg, M.-L., Buck, S.H. and Lovenberg, W., Development of an anti-idiotypic antibody that blocks substance P primary antibodies and substance P membrane binding, 131

Tanimoto, M. and Okada, Y., The protective effect of hypothermia on hippocampal slices from guinea pig during deprivation of oxygen and glucose, 239
Tavares, M.A., see Paula-Barbosa, M., 139
Terashima, T., Katada, T., Oinuma, M., Inoue, Y. and Ui, M., Endocrine cells in pancreatic islets of Langerhans are immunoreactive to antibody against guanine nucleotide-binding protein (G_o) purified from rat brain, 190
Thach, W.T., see Mink, J.W., 393
Thierry, A.M., see Ferino, F., 257
Thomas, H.C. and Espinoza, S.G., Relationships between interhemispheric cortical connections and visual areas in hooded rats, 214
Thompson, G.C., see Usami, S., 367
Tokutomi, N., see Oyama, Y., 143
Townsend, C.M., see Hsu, L.L., 232
Toyka, K.V., see Hartung, H.-P., 347
Tsin, A.T.C., see Vaughan, M.K., 321

Ui, M., see Terashima, T., 190
Upp Jr., J.R., see Hsu, L.L., 232
Usami, S., Igarashi, M. and Thompson, G.C., GABA-like immunoreactivity in the squirrel monkey vestibular endorgans, 367

Vaněček, J., see Illnerová, H., 367
Vaughan, G.M., see Vaughan, M.K., 321
Vaughan, M.K., Chambers, J.P., Tsin, A.T.C., Vaughan, G.M. and Reiter, R.J., Pineal and retinal lysosomal enzyme rhythms, 321
Vezina, P., Kalivas, P.W. and Stewart, J., Sensitization occurs to the locomotor effects of morphine and the specific μ opioid receptor agonist, DAGO, administered repeatedly to the ventral tegmental area but not to the nucleus accumbens, 51

Wahlestedt, C., Skagerberg, G., Ekman, R., Heilig, M., Sundler, F. and Håkanson, R., Neuropeptide Y (NPY) in the area of the hypothalamic paraventricular nucleus activates the pituitary-adrenocortical axis in the rat, 33
Wainer, B.H., see Mufson, E.J., 385
Wang, M.R., see Kuo, J.S., 181
Wilson, D.A. and Leon, M., Evidence of lateral synaptic interactions in olfactory bulb output cell responses to odors, 175
Wirsig, C.R. and Leonard, C.M., Terminal nerve damage impairs the mating behavior of the male hamster, 293
Wright, E.M., see Saito, Y., 267

Yashima, Y., see Nakajima, T., 360
Yu, C.Y., see Kuo, J.S., 181
Yu, J.-R., see Hsu, L.L., 232

Zill, S.N., Selective mechanical stimulation of an identified proprioceptor in freely moving locusts: role of resistance reflexes in active posture, 195